

## Original Article

# Efficiency of Commonly Used Disinfectants and Antiseptics against Pathogenic Bacteria Isolated from Tripoli University Hospital, Libya

Mahmoud Ashawesh<sup>1</sup>\*, Sara Alashhab, Sakena Kori

Department of Medical Laboratories Sciences, Faculty of Medical Technology, the University of Tripoli, Libya

Corresponding Email: [M.ashawesh@uot.edu.ly](mailto:M.ashawesh@uot.edu.ly)**ABSTRACT**

**Background and objectives:** Disinfectants and antiseptics are widely used in hospitals and other healthcare centers to ensure the prevention of nosocomial infections and provide efficient infection control. Numerous studies have recently shown that hospital-acquired bacteria are developing resistance against such compounds. The purpose of this study is to examine the effectiveness of the commonly used disinfectants and antiseptics against a group of nosocomial bacteria isolated from Tripoli University Hospital, Libya. **Material and Methods:** Five bacterial pathogens isolated from different hospital departments are *klebsiella* spp., *Proteus* spp., *Escherichia coli* (*E. coli*), *Acinetobacter* spp., *Staphylococcus aureus* (*S. aureus*). These isolated bacterial species were subjected to disinfectants and antiseptics impregnated with filter paper discs (disk diffusion assay). The disinfectants and antiseptics used were; Hydrogen peroxide ( $H_2O_2$ ), OROLIN® Multisept plus (Orolin), Chlorhexidine (CHX), 84 Disinfectant, ACTOSAL® Flache AF (Actosal) and Ethyl alcohol. **Results:** In the present study found that the most effective was  $H_2O_2$  antiseptic against the tested bacterial pathogens with inhibition diameters of 10 to 26mm, except for *Proteus* bacteria which illustrated resistance at high concentrations. Similarly, Orolin disinfectant shows very excellent efficacy against tested bacteria, and even with low concentrations. Good efficacy was observed for Actosal disinfectant on all species with diameters of inhibition ranging from 10 to 15mm. *S. aureus* in particular was very sensitive to 70% Ethyl alcohol (20mm), while all tested species were slightly insensitive to both Ethyl alcohol and CHX. In contrast, no effect of the 84 disinfectant was observed on most of isolates. **Conclusion:** The results of this study indicated that Orolin disinfectant had comparable effects to  $H_2O_2$  atiseptic but less than to Actosal, 70% Ethyl alcohol and CHX, whereas 84 disinfectant was the weakest one. This study suggests the necessity of applying continuous monitoring to determine the antimicrobial efficacy of these antimicrobial agents regularly.

**Keywords:** Disinfectants, Antiseptics, Tolerance, Resistance, Nosocomial Infections.**Citation:** Ashawesh M, Alashhab S, Kori S. Efficiency of Commonly Used Disinfectants and Antiseptics against Pathogenic Bacteria Isolated from Tripoli University Hospital, Libya. *Khalij-Libya J Dent Med Res.* 2024;8(1):119–128.<https://doi.org/10.47705/kjdmr.2481017>**Received:** 12/04/24; **accepted:** 07/06/24; **published:** 11/06/24Copyright © Khalij-Libya Journal (KJDMR) 2024. Open Access. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>

**الخلفية والأهداف:** تستخدم المطهرات والمعقمات على نطاق واسع في المستشفيات ومراكز الرعاية الصحية لضمان الوقاية من عدوى المستشفيات وتوفير مكافحة العدوى بكفاءة. أظهرت العديد من الدراسات مؤخرًا أن البكتيريا المكتسبة من المستشفيات تطور مقاومة ضد هذه المركبات. الغرض من هذه الدراسة هو فحص مدى فعالية المطهرات والمعقمات الشائعة الاستخدام ضد مجموعة من البكتيريا المعزولة من مستشفى جامعة طرابلس، ليبيا. **طرق الدراسة:** تم الحصول على خمسة أنواع بكتيرية معزولة من مختلف أقسام المستشفى الجامعي وهي *klebsiella* spp., *Proteus* spp., *E. coli*, *Acinetobacter* spp., *S. aureus*. أجريت زراعة هذه الأنواع البكتيرية المعزولة في مولر هينتون آجار (MHA) و تعرضت للمطهرات والمعقمات المشربة بأقراص ورق الترشيح. وكانت المطهرات والمعقمات المستخدمة هي: بيروكسيد الهيدروجين ( $H_2O_2$ )، OROLIN® Multisept plus (الأورولين)، الكلوروكسيدين (CHX)، مطهر 84، ACTOSAL® Flache AF (أكتوسال) والكحول الإيثيلي. **النتائج:** في الدراسة الحالية وجد أن المطهر الأكثر فعالية هو  $H_2O_2$  ضد البكتيريا المعزولة التي تم اختبارها بأقراص تنبسط تراوحت بين 10 إلى 26 ملم، باستثناء بكتيريا *Proteus* التي أظهرت مقاومة عند التراكيز العالية. وبالمثل، يُظهر مطهر الأورولين فعالية ممتازة جدًا ضد البكتيريا التي تم اختبارها، وحتى مع التركيزات المنخفضة. وقد لوحظت فعالية جيدة لمطهر أكتوسال على جميع الأنواع التي تتراوح أقطار تنبسطها من 10 إلى 15 ملم. كانت *S. aureus* على وجه الخصوص حساسة جدًا للكحول الإيثيلي (20 مم)، في حين كانت جميع الأنواع التي تم اختبارها قليلة الحساسية لكل من الكحول الإيثيلي و CHX. في المقابل، لم يلاحظ أي تأثير للمطهر 84 على أغلب البكتيريا المعزولة. **الخلاصة:** أشارت نتائج هذه الدراسة إلى أن مطهر الأورولين كان له تأثيرات مماثلة لمطهر  $H_2O_2$  ولكن أقل من أكتوسال والكحول الإيثيلي و CHX، في حين كان مطهر 84 هو الأضعف. تشير هذه الدراسة إلى ضرورة تطبيق المراقبة المستمرة لتحديد فعالية مضادات الميكروبات لهذه العوامل المضادة للميكروبات بانتظام.

## INTRODUCTION

Antibiotic-resistant bacteria are becoming more common sources of hospital-acquired infections. This has led to high rates of morbidity and mortality among patients, which significantly raises the cost of care due to longer hospital stays and the requirement for more expensive medications [2,1]. For instance, methicillin-resistant *Staphylococcus aureus* (MRSA) is the main cause of nosocomial infections that are becoming increasingly difficult to combat due to developed resistance to all classes of present antibiotics [3]. It has been postulated that countries with low incidence of MRSA infections tend to use stronger antibiotic restriction policies and implement more stringent infection control protocols [2,4]. Nowadays, novel antimicrobial agents are desperately needed because those currently available have become ineffective due to the emergence of antibiotic resistance in hospitals and communities [5]. One way to eliminate the global spread of antibiotic resistance and control the circulation of resistant bacteria in hospitals, homes and communities is the constant use of disinfectants and antiseptics [6]. These substances are antimicrobial slowing or killing germs whether if presented on a non-living object (disinfectants) or on the living tissues (antiseptics) and the main reason for using such substances is to reduce exposure to infection. However, with longer use the germs might also have acquired resistance to disinfectants of various kinds [7]. In recent years, numerous publications have demonstrated that bacteria can develop resistance not only to antibiotics but also tolerance to sanitizers and disinfectants [8-10]. In fact, these substances are widely used in hospitals and other healthcare centers to control microbial growth in living tissues and non-living objects and are considered an essential part of infection control practices and help prevent hospital infections [11].

Alcohols are one of the most important sanitizing antimicrobial agents that can denature proteins, meaning that, gives excellent bactericidal efficacy. Alcohol is essential in medical detergents and

disinfectants and becoming an alternative to hand washing in healthcare settings and some public places due to its effectiveness in the rapid destruction of some pathogens [12]. Remarkably, substantial evidence exists supporting the emergence of antimicrobial-resistant bacterial isolates. For example, a study conducted by Pidot *et al.*, 2018 showed that *Enterococcus faecium* isolates were tolerant to isopropanol alcohol [13]. Furthermore, Akin and co-workers have detected resistant isolates against 3% H<sub>2</sub>O<sub>2</sub> disinfectant [14]. In addition, a recent study by Morante *et al.*, 2021 elucidated resistance to CHX and isopropanol alcohol among *Klebsiella pneumoniae* isolates found in Peru hospitals [15]. Thus, over the past few years, there has been an increased concern due to the misuse of these disinfectants and antiseptics worldwide which as a consequence might cause bacteria to develop resistance [16].

Currently, a new concern from alcohol-tolerant bacteria has started to be developed [17]. In a previous study, researchers examined the alcohol tolerance of 139 hospital *E. faecium* isolates collected from 1997 to 2015 from two Melbourne hospitals and examined how well these isolates survived when treated with dilute isopropyl alcohol. Interestingly, they found that samples collected after 2010 were more tolerant to 70% isopropanol surface disinfection compared to isolates collected before 2010 [13]. Similarly, another study demonstrated the inhibitory effects of five common antiseptics; CHX, H<sub>2</sub>O<sub>2</sub>, Iodine, Ethanol, and Dettol on MRSA, *Acinetobacter baumannii*, *E. coli*, and *klebsiella* species, and significantly found that all tested bacteria were resistant to ethanol at all concentrations. Indeed, this study also showed that H<sub>2</sub>O<sub>2</sub> was the most effective disinfectant compared to other disinfectants followed by CHX [18].

It could be argued that either due to overuse or misuse of disinfectants and consumption of low-quality products, tolerance or even resistance against these disinfectants can occur [19, 20]. Consequently, as a result of disinfectants possibly being no longer as effective as should be, this would seriously

threaten clinical settings and patient safety [21]. The aim of this study is to examine the effectiveness of commonly used disinfectants and antiseptics in Tripoli university hospital against the most common nosocomial bacteria and determine if the used concentrations at these healthcare facilities are capable or not of eliminating the superbug.

## METHODS

### *Microbial strains*

Five strains of pathogenic bacteria were used, of which one was Gram-positive *Staphylococcus aureus* (*S. aureus*), and four were Gram-negative *klebsiella* spp., *Proteus* spp., *Escherichia coli* (*E. coli*), *Acinetobacter* spp. All bacteria were obtained from the different departments at Tripoli University Hospital and identified by standard microbiological tests. These isolates were used to evaluate the inhibitory activity of disinfectants and antiseptics against them.

### *Culture preparation and susceptibility test*

A swab was taken from a colony of bacteria using a sterile cotton swab dipped in normal saline, and the swab was distributed over the entire Muller-Hinton dish. Cultivate bacteria was tested for the effect of antiseptics and sterilizers using the (Disc Diffusion Method). The selected disinfectants or antiseptics were impregnated with filter paper discs with a diameter of (6mm) saturated with the specific concentrations and used separately. Then sterile forceps with a pointed end were used to transfer the discs to the surface of the implanted dish and lightly pressed the surface to fix it on the culture medium. To avoid overlapping the areas of inhibition, an appropriate distance was left between one disc and another. Then all dishes were placed in the incubator for 24 hours at 37°C. After the incubation period, the diameter of the bacterial growth inhibition zones was measured around the discs, which appear as a halo without bacterial growth surrounding the disinfectant disc.

### *Disinfectants and antiseptics*

Hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>), OROLIN® Multisept plus (Orolin), Chlorhexidine (CHX), 84 Disinfectant, ACTOSAL® Flache AF (Actosal) and Ethyl alcohol were used in this study to test their efficiency in eliminating bacteria and examine the sensitivity of bacteria to such compounds. These antimicrobial substances are commonly used in Tripoli University Hospital and were prepared according to manufacturer recommendations. The manufacturer and other information of such substances are shown in Table 1.

As recommended by the manufacturer, several increasing effective concentrations of these substances were prepared using the dilution methods to test in this study. Furthermore, some particular concentrations recommended by the hospital were also examined (Table 1). A dilution procedure was done (1:100) where the original concentration of the substance was considered 100%, and the subsequent concentration was prepared by adding 1 ml of disinfectant in a tube with 99 ml of distilled water to give a concentration of 1% and so on. Depending on the type of disinfectants or antiseptics used in this study, different concentrations were applied in this study (Table 1).

Table 1: Types and concentration of disinfectants and antiseptics used in the study.

Trade name	R/Con.	H/Con.	Main effective ingredient	Manufacture
Hydrogen Peroxide (H <sub>2</sub> O <sub>2</sub> )	3%, 6%	3%, 6%	Hydrogen Peroxide (H <sub>2</sub> O <sub>2</sub> )	(AQUA) Turkey
OROLIN® Multisept plus	2%	2%, 5%, 10%, 15%	<ul style="list-style-type: none"> <li>Didecyldimethylammonium chloride &gt;2.5% – &lt;5%</li> <li>N-(3-Aminopropyl)-N-dodecylpropane -1,3-diamine &gt;5% – &lt;15%</li> <li>2-amino-2-methylpropanol &gt;2.5% – &lt;5%</li> </ul>	(DENTAID) Spain
Chlorhexidine (CHX)	100%	100%	<ul style="list-style-type: none"> <li>Chlorhexidine 0.12%</li> <li>Cetylpyridinium chloride 0.05%</li> </ul>	(DENTAID) Spain
84 Disinfectant	1%	1%, 2%, 5%	Parachlormetaxylenol 4.5% - 6.5%	(GIG) Germany
ACTOSAL® Flache AF	100%	100%	<ul style="list-style-type: none"> <li>Poly(oxy-1,2-ethanediyl), alpha-(2-propylheptyl)-omegahydroxy &gt; 9% – &lt; 11%</li> <li>Didecyldimethylammonium chloride &gt; 7% – &lt; 9%</li> <li>Polyhexamethylene Biguanide Hydrochlorid &gt; 2% – &lt; 3%</li> </ul>	(ACTO) Germany
ETHANOL	70%	70%	Ethyl alcohol	(WAN CARE) Turkey

## RESULTS

In order to examine the efficiency of using different concentrations of Orolin disinfectant as recommended by Tripoli university hospital (Table 1) against our bacteria isolates 2%, 5%, 10%, and 15% of disinfectant concentrations were prepared and impregnated into paper discs and exposed to bacterial isolates as described in material and methods. As shown in Figure 1, Table 2 and Figure 3A, our results of the disc diffusion method showed a sensitivity of tested microorganisms to disinfectant Orolin at different concentrations which were compared to the antiseptic alcohol at a concentration of 95% as a control. These results revealed a clear inhibitory activity of Orolin disinfectant at selected concentrations against all types of bacteria and were apparent at the highest concentration (15%) for most of the bacterial strains suggesting that all isolates exhibited excellent sensitivity to Orolin. The sensitivity to 95% alcohol was very limited due to alcohol evaporation over time, (Figure 1).

To test the growth rate of our bacterial isolates against all disinfectants and antiseptics listed in

Table 1, recommended concentrations whether from the hospital or from the manufacturer were applied in this study. The results showed that the five types of bacteria isolates (*Klebsiella* spp., *Proteus* spp., *E. coli*, *Acinetobacter* spp., and *S. aureus*) were sensitive to H<sub>2</sub>O<sub>2</sub> at both used concentrations (3% & 6%) with inhibition diameters ranging between 10-26mm except for the *Proteus* spp. which shows resistance at 6% concentration (Table 2 and Figure 3C). Interestingly, despite *E. coli* bacteria having clear sensitivity to most disinfectants, it shows a clear resistance to 84 disinfectants at all given concentrations (Table 2 and Figure 3B). Similarly, *Acinetobacter* spp., *Proteus* spp. illustrated resistance to 84 disinfectants at 2% and 5% concentration (Table 2 and Figure 3B), suggesting a possible emerging resistance against 84 Disinfectant. Although *S. aureus* demonstrated tolerance against 84 disinfectants at concentration of 2% and 5% with small zone of




Figure 1: Inhibition zones caused by Orolin at selected concentrations of 5%,10% as well as 15% and alcohol at 95% on the studied bacteria.

Table 2: The distribution of bacteria sensitivity against disinfectant concentrations, the numbers represent the diameters of the inhibition zone (mm).

		Antiseptics and disinfectant concentrations											
		Orolin 2%	Orolin 5%	Orolin 10%	Orolin 15%	84 D 1%	84 D 2%	84 D 5%	Ethyl 70%	CHX	Acto AF	H <sub>2</sub> O <sub>2</sub> 3%	H <sub>2</sub> O <sub>2</sub> 6%
Tested Bacteria	Acinetobacter	16	12	18	22	6.5	R	R	8	9	10	15	18
	Proteus	16	17	20	20	6.5	R	R	7	6.5	12	10	R
	S. aureus	22	20	20	20	R	8	9	20	13	14.5	26	18
	E. coli	14	17	19	20	R	R	R	14	9	15	18	20
	Klebsiella	10	22	20	25	6.5	6.5	6.5	10	12	15	17	18

Oro = Orolin Multisept Plus / 84 = 84 Disinfectant / Ethyl = Ethyl alcohol / CHX = Chlorhexidine / Acto = Actosal Flache AF / H<sub>2</sub>O<sub>2</sub> = Hydrogen Peroxide. The yellow shading indicates resistance. Grey shading shows maximum sensitivity.

inhibition ranging between 8 to 9mm, its resistance at the lowest concentration was detectable (Table 2). On the other hand, *Klebsiella* bacteria were more sensitive to all disinfectants and recorded zone of inhibition ranged between 6.5-25mm.

Orolin disinfectant was very effective against all bacteria tested and its inhibitory effect was increasing at higher concentrations followed by H<sub>2</sub>O<sub>2</sub>. Conversely, CHX had comparable effects to Actosal but less than ethyl alcohol (Table 2). *S. aureus* species in particular represented a maximum zone of inhibition against 3% H<sub>2</sub>O<sub>2</sub> (26mm) compared to the other tested disinfectants (Table 2 and Figure 2). In support of these results, it is clear as shown in Figure 3 that 84 Disinfectant was the least effective for all types of bacteria, followed by 70% Ethyl alcohol and CHX. Our results showed that Orolin was the most effective disinfectant than the others followed by H<sub>2</sub>O<sub>2</sub> (Figure 3A and C).



Figure 2: Inhibition growth zone test of disinfectants and antiseptics used in the study toward *S. aureus*. H<sub>2</sub>O<sub>2</sub> 3% [1], 10% Orolin [2], 100% CHX [3], 84 Disinfectant 1% [4], 100% Actosal [5] and 70% Ethyl alcohol [6].




Figure 3: Distribution of bacteria sensitivity against disinfectant concentrations. A is the distribution of bacteria sensitivity against Orolin and 84 Disinfectant disinfectants. B is the distribution of bacteria sensitivity against Ethyl alcohol. C is for CHX, Actosal and H<sub>2</sub>O<sub>2</sub>.

## DISSCUSSION

The isolated bacteria obtained in this study were subjected to susceptibility tests and exposed to several disinfectants and antiseptics used to control microbial infection in Tripoli University hospitals. The purpose of this test was to evaluate the inhibitory effect of these antimicrobial compounds.

Clearly, the five tested types of bacteria used (*Klebsiella* spp., *Proteus* spp., *E. coli*, *Acinetobacter* spp., and *S. aureus*) were sensitive to the  $H_2O_2$  at a concentration of (3% and 6%) produced zone of inhibition, ranging between 10 mm to 26mm, except that *Proteus* bacteria was capable to resist this compound at a concentration of 6% (Table 2). In agreement with this, an observation was conducted by Hassanain *et al.*, 2019, concluding that  $H_2O_2$  was the most effective antiseptic used in the study as demonstrated excellent inhibitory effects on all tested nosocomial bacteria even with lower concentrations [18]. Similarly, in support of this finding Lineback and co-workers found that  $H_2O_2$  antiseptic had significantly higher bactericidal efficacies when compared with other disinfectants [22]. It was envisaged that this action of  $H_2O_2$  is due to a potent oxidant that produces hydroxyl radicals which as a consequence will perturb the cell membrane, lipids and other essential cell components [18,23].

On the other hand, Orolin disinfectant was shown to be the strongest disinfectant used, and even that at the lowest concentrations it still had excellent inhibitory effects among tested bacteria (Table 2). Orolin is a powerful disinfectant due to its distinctive composition includes strong corrosion inhibitors [24]. Obviously, Orolin gives inhibitory zones ranged between (10-22mm) at 2% concentration and increased at 15% concentration to reach between (20-25mm), which demonstrates a growth decline in a disinfectant dose-dependent manner (Table 2, Figures 1 and Figure 3A).

Unlike Orolin disinfectant, 84 disinfectants did not produce any significant zones of inhibition against the tested organisms even at increasing concentrations (Table 2). In fact, 84 disinfectant was ineffective against the tested bacteria with comparable zone sizes to other used disinfectants (Figure 3B). Surprisingly, the most tested bacteria (*Acinetobacter* spp, *Proteus* spp. and *E. coli*) were resistant to this particular disinfectant and both *S. aureus* and *Klebsiella* spp. illustrated tolerance, suggesting that the manufacturer should take this

into consideration (Table 2 Figure 2). 84 disinfectant is sodium hypochlorite containing 5.5% to 6.5% chlorine and resistance against this compound was indeed documented [25, 26]. It was thus predicted that the dramatic increase in the use of this disinfectant during the COVID-19 epidemic has a key role in developing bacterial resistance in clinical settings [25].

Actosal is widely used as a surface disinfectant in hospitals and contains mainly diethyldimethylammonium chloride (quaternary ammonium compound commonly applied for cleaning and disinfection purposes) [27]. In this study, Actosal demonstrated decent efficacy against the clinical isolates with diameters of inhibition ranging from 10 to 15mm (Table 2), although reduced susceptibility to diethyl dimethylammonium chloride was reported [16,22,28]. Alternatively, 70% Ethyl alcohol revealed a quite variation in zones of inhibition against the tested organisms. We noticed that it showed good activity against *S. aureus* (20mm) indicating susceptibility, while a moderate effect was evident in both *E. coli* and *Klebsiella* bacteria with diameters ranging between (10-14mm) (Table 2). Nevertheless, both *Proteus* and *Acinetobacter* species were Ethyl alcohol-tolerant bacteria (Table 2). Finally, a lower level of growth activity was detected in the studied bacteria when treated with the oral CHX antiseptic which exhibited a zone of inhibition in the range of 6.5-13mm (Table 2 and Figure 3C). CHX is also used in hospital settings such as in cleaning surfaces and surgical sites but is mainly applied in dental practice. Interestingly, numerous recent publications have indicated the emergence of resistance to CHX [15,29,30,31]. In general, the results of this study indicated that  $H_2O_2$  had comparable effects to Orolin disinfectant but less than to Actosal, Ethyl alcohol and CHX.

While there is a good understanding of antibiotic resistance, tolerance to disinfectants remains unclear and yet under investigation. Unlike antibiotics, disinfectants mode of action can be nonspecific, targeting different mechanisms or sites in bacterial

cells which is subsequently imposing cell disruption. Thereby, conferring resistance is unlikely [32,33]. Conversely, tolerance or even resistance against overuse or misuse disinfectants can be acquired by a number of mechanisms such as; (i) developing mutation of an endogenous chromosomal gene, (ii) incorporating mobile genetic elements such as plasmids, jumping transposons and other cell elements, and (iii) due to changes in cell membrane permeability or increased efflux pump expression [16,20]. In addition, a threaten linkage between long term usage of disinfectants and emergence of resistance to antibiotics was recently reported [34]. This is particularly a matter of concern since available data on this situation here in Libya is scarce. At present, regulations for antibiotic consumption worldwide are stringent. Disinfectants are however subject to fewer restrictions compared to antibiotics. Therefore, it is imperative to call for the controlled use of disinfectants to manage the rise of antibiotic resistance caused by prolonged exposure to these substances [34].

## CONCLUSION

This study confirms the efficient ability of both Orolin disinfectant and H<sub>2</sub>O<sub>2</sub> antiseptic to degrade all isolated bacterial species followed by Actosal and 70% Ethyl alcohol. However, 84 disinfectant was the weakest one and showed no significant effects on these species. Collectively, this evidence suggests the urgent need for continuous surveillance to determine the antimicrobial efficiency of disinfectants and antiseptics regularly. We recommend a daily review of the effectiveness of disinfectants and chemical sterilizers against multidrug-resistant pathogens is deemed necessary. Further comprehensive studies are required to investigate this issue by collecting larger samples from different hospitals and healthcare centers and using a broad collection of antimicrobial agents to determine whether the widespread use of these products is responsible for the development of microbial resistance or not.

## Conflict of Interest

There are no financial, personal, or professional conflicts of interest to declare.

## REFERENCES

1. Emori TG, Gaynes RP. An overview of nosocomial infections, including the role of the microbiology laboratory. *Clin. Microbiol. Rev.* 1993 Oct;6(4):428-42.
2. Struelens MJ. The epidemiology of antimicrobial resistance in hospital acquired infections: problems and possible solutions. *BMJ.* 1998;317(7159):652-4.
3. Kourtis AP, Hatfield K, Baggs J, Mu Y, See I, Epsom E, Nadle J, Kainer MA, Dumyati G, Petit S, Ray SM. Emerging Infections Program MRSA author group; Ham D, Capers C, Ewing H, Coffin N, McDonald LC, Jernigan J, Cardo D. Vital Signs: Epidemiology and Recent Trends in Methicillin-Resistant and in Methicillin-Susceptible *Staphylococcus aureus* Bloodstream Infections - United States. *MMWR Morb. Mortal. Wkly. Rep.* 2019 Mar 8;68(9):214-219.
4. Kinoshita T, Tokumasu H, Tanaka S, Kramer A, Kawakami K. Policy implementation for methicillin-resistant *Staphylococcus aureus* in seven European countries: a comparative analysis from 1999 to 2015. *J. Mark Access Health Policy.* 2017 Jul 26;5(1):1351293.
5. Huttner A, Harbarth S, Carlet J, Cosgrove S, Goossens H, Holmes A, Jarlier V, Voss A, Pittet D. Antimicrobial resistance: a global view from the 2013 World Healthcare-Associated Infections Forum. *Antimicrob. Resist. Infect. Control.* 2013 Nov 18;2:31.
6. Maillard JY, Bloomfield SF, Courvalin P, Essack SY, Gandra S, Gerba CP, Rubino JR, Scott EA. Reducing antibiotic prescribing and addressing the global problem of antibiotic resistance by targeted hygiene in the home and everyday life settings: A position paper. *Am. J. Infect. Control.* 2020 Sep;48(9):1090-1099.
7. Fernandes ÂR, Rodrigues AG, Cobrado L. Effect of prolonged exposure to disinfectants in the antimicrobial resistance profile of relevant microorganisms: a systematic review. *J. Hosp. Infect.* 2024 May 11:S0195-6701(24)00163-4.
8. Yeung YS, Ma Y, Liu SY, Pun WH, Chua SL. Prevalence of alcohol-tolerant and antibiotic-resistant bacterial pathogens on public hand



- sanitizer dispensers. *J Hosp Infect.* 2022 Sep;127:26-33.
9. Lim K, Li WY, Dinata A, Ho ET. Comparing the antibacterial efficacy and functionality of different commercial alcohol-based sanitizers. *PLoS. One.* 2023 Mar 27;18(3):e0282005.
  10. Chojnacki M, Dobrotka C, Osborn R, Johnson W, Young M, Meyer B, Laskey E, Wozniak RAF, Dewhurst S, Dunman PM. Evaluating the Antimicrobial Properties of Commercial Hand Sanitizers. *mSphere.* 2021 Mar 3;6(2):e00062-21.
  11. Saha, A., Haque, M., Karmaker, S., & Mohanta, M. Antibacterial Effects of Some Antiseptics and Disinfectants. *Journal of Life and Earth Science.* 2011; 3, 19–21.
  12. Todd EC, Michaels BS, Holah J, Smith D, Greig JD, Bartleson CA. Outbreaks where food workers have been implicated in the spread of foodborne disease. Part 10. Alcohol-based antiseptics for hand disinfection and a comparison of their effectiveness with soaps. *J. Food Prot.* 2010 Nov;73(11):2128-40.
  13. Pidot SJ, Gao W, Buultjens AH, Monk IR, Guerillot R, Carter GP, Lee JYH, Lam MMC, Grayson ML, Ballard SA, Mahony AA, Grabsch EA, Kotsanas D, Korman TM, Coombs GW, Robinson JO, Gonçalves da Silva A, Seemann T, Howden BP, Johnson PDR, Stinear TP. Increasing tolerance of hospital *Enterococcus faecium* to handwash alcohols. *Sci. Transl. Med.* 2018 Aug 1;10(452):eaar6115.
  14. Eryilmaz M, Akin A, Arıkan Akan O. Investigation of the efficacy of some disinfectants against nosocomial *Staphylococcus aureus* and *Enterococcus* spp. isolates. *Mikrobiyol Bul.* 2011 Jul;45(3):454-60.
  15. Morante J, Quispe AM, Ymaña B, Moya-Salazar J, Luque N, Soza G, Ramos Chirinos M, Pons MJ. Tolerance to disinfectants (chlorhexidine and isopropanol) and its association with antibiotic resistance in clinically-related *Klebsiella pneumoniae* isolates. *Pathog. Glob. Health.* 2021 Feb;115(1):53-60.
  16. Boyce JM. Quaternary ammonium disinfectants and antiseptics: tolerance, resistance and potential impact on antibiotic resistance. *Antimicrob. Resist. Infect. Control.* 2023 Apr 13;12(1):32.
  17. Yeung W, Ma Y., Liu Y, Pun H., Chua .L. Prevalence of alcohol -tolerant and antibiotic-resistant bacterial pathogens on public hand sanitizer dispensers, *Journal of Hospital Infection.* 2022;127: 26-33.
  18. Hassanain T., Alyaa, A., Ahmad, R., Nadia Z., Syakirah H., Nur M. & Amilah K. Effectiveness of commonly used antiseptics on bacteria causing nosocomial infections in tertiary hospital in Malaysia. *African Journal of Microbiology Research.* 2019;13(10), 188-194.
  19. Hamad, B. Study in The Inhibitory Effect of Disinfectants and Antiseptic Traded From Local Market in Nineveh The Province of on Some Pathogenic Bacteria. *Journal of Education and Science,* 2019; 28(2): 103-112.
  20. Rozman U, Pušnik M, Kmetec S, Duh D, Šostar Turk S. Reduced Susceptibility and Increased Resistance of Bacteria against Disinfectants: A Systematic Review. *Microorganisms.* 2021 Dec 10;9(12):2550.
  21. Tong C, Hu H, Chen G, Li Z, Li A, Zhang J. Disinfectant resistance in bacteria: Mechanisms, spread, and resolution strategies. *Environ. Res.* 2021 Apr;195:110897.
  22. Lineback B., Nkemngong A., Wu T. et al. Hydrogen peroxide and sodium hypochlorite disinfectants are more effective against *Staphylococcus aureus* and *Pseudomonas aeruginosa* biofilms than quaternary ammonium compounds. *Antimicrob. Resist. Infect. Control.* 2018; 7, 154.
  23. Mai-Prochnow A, Lucas-Elio P, Egan S, Thomas T, Webb JS, Sanchez-Amat A, Kjelleberg S. Hydrogen peroxide linked to lysine oxidase activity facilitates biofilm differentiation and dispersal in several gram-negative bacteria. *J. Bacteriol.* 2008 Aug;190(15):5493-501.
  24. [https://oroclean.com/ch-en/products/instrument-disinfectants/OROLIN\\_Multisept\\_Plus](https://oroclean.com/ch-en/products/instrument-disinfectants/OROLIN_Multisept_Plus)
  25. Jin M, Liu L, Wang DN, Yang D, Liu WL, Yin J, Yang ZW, Wang HR, Qiu ZG, Shen ZQ, Shi DY, Li HB, Guo JH, Li JW. Chlorine disinfection promotes the exchange of antibiotic resistance genes across bacterial genera by natural transformation. *ISME J.* 2020 Jul;14(7):1847-1856.
  26. Sultana S, Foti A, Dahl JU. Bacterial Defense Systems against the Neutrophilic Oxidant Hypochlorous Acid. *Infect. Immun.* 2020 Jun 22;88(7):e00964-19.
  27. <https://actomarkt.com/en/products/actosal-flache-af>
  28. Soumet C, Méheust D, Pissavin C, Le Grandois P, Frémaux B, Feurer C, Le Roux A, Denis M, Maris P. Reduced susceptibilities to biocides and resistance to antibiotics in food-associated bacteria following

- exposure to quaternary ammonium compounds. *J. Appl. Microbiol.* 2016 Nov;121(5):1275-1281.
29. Roode GJ, Bütow KW. A Descriptive Study of Chlorhexidine as a Disinfectant in Cleft Palate Surgery. *Clin. Med. Res.* 2018 Jun;16(1-2):9-15.
30. Abbood M, Hijazi K, Gould M. Chlorhexidine Resistance or Cross-Resistance, That Is the Question. *Antibiotics (Basel)*. 2023 Apr 22;12(5):798.
31. Cieplik F, Jakubovics NS, Buchalla W, Maisch T, Hellwig E, Al-Ahmad A. Resistance Toward Chlorhexidine in Oral Bacteria - Is There Cause for Concern? *Front Microbiol.* 2019 Mar 22;10:587
32. Goudarzi, M. and Navidinia, M. Overview perspective of bacterial strategies of resistance to biocides and antibiotics. *Arch. Clin. Infect. Dis.* 2019, 14, e65744.
33. Romero JL, Grande Burgos MJ, Pérez-Pulido R, Gálvez A, Lucas R. Resistance to Antibiotics, Biocides, Preservatives and Metals in Bacteria Isolated from Seafoods: Co-Selection of Strains Resistant or Tolerant to Different Classes of Compounds. *Front Microbiol.* 2017 Aug 31;8:1650.
34. Wu-Chen RA, Feng J, Elhadidy M, Nambiar RB, Liao X, Yue M, Ding T. Long-term exposure to food-grade disinfectants causes cross-resistance to antibiotics in *Salmonella enterica* serovar Typhimurium strains with different antibiograms and sequence types. *Antimicrob. Resist. Infect. Control.* 2023 Dec 13;12(1):145.
35. Wu RA, Feng J, Yue M, Liu D, Ding T. Overuse of food-grade disinfectants threatens a global spread of antimicrobial-resistant bacteria. *Crit Rev Food Sci. Nutr.* 2023 Feb 9:1-10.